

Standard report for Vivax Malaria

WWARN Vivax Primaquine Study Group

For further information go to <https://www.iddo.org/wwarn/vivax-reports>

09 October, 2025

Introduction

This report has been produced for countries: Thailand

The studies included within this report are shown in Table 0.

Table 0: Studies included in this report

Author-year	Country	Recruitment Period	Age range (years)	Follow up (days)	Included treatment arms*	PQ supervision	Patients available
Chu-2018	Thailand	2010 - 2011	1.5 - 63	365	Cq_Pq_7.0_14d_D0, Cq	Fully supervised	420
Longley-2016	Thailand	2014	7 - 71	270	Cq_Pq_3.5_14d_D1	Fully supervised	43
Llanos-Cuentas-2019	Thailand	2015 - 2016	15 - 48	180	Cq_Pq_3.5_14d_D1	<50% supervised	8
Lacerda-2019	Thailand	2014 - 2017	18 - 49	180	Cq_Pq_3.5_14d_D1, Cq	<50% supervised	30
Llanos-Cuentas-2014	Thailand	2011 - 2013	18 - 58	180	Cq, Cq_Pq_3.5_14d_D1	<50% supervised	32
Chu-2019	Thailand	2012 - 2014	1.5 - 63	365	DP_Pq_7.0_7d_D0, DP_Pq_7.0_14d_D0, Cq_Pq_7.0_7d_D0, Cq_Pq_7.0_14d_D0	Fully supervised	654
Pukrittayakamee-2010	Thailand	1995 - 1998	14 - 61	28	Pq_7.0_7d_D0, Pq_3.5_7d_D0	Fully supervised	85

* ACT – artemisinin-based combination treatment; As – artesunate; AL – artemether -lumefantrine; Aq – amodiaquine; Cq – chloroquine; DP – dihydroartemisinin-piperaquine; GI – gastrointestinal; Mf – mefloquine; PQ/Pq – primaquine; SP – sulfadoxine-pyrimethamine;

Treatment code describes (schizontocidal drug)(hypnozoitocidal drug)(total primaquine dose)(duration of primaquine treatment eg 14d = 14 days)(primaquine start day)

1: EFFICACY

1.1: Description

The efficacy study was undertaken to better understand the impact of primaquine dose on the prevention of *P. vivax* recurrences. Inclusion in the efficacy meta-analysis was restricted to studies with 42 days or more follow up and patients with data on day 0 parasitaemia.

In this report the efficacy study includes 1187 patients across 8 study sites, from 6 studies.

1.2: Characteristics of Study Population

Table 1_eff: Characteristics of the study population for the efficacy study analysis, categorised by total primaquine category

	Primaquine (PQ) Treatment				Total (N=1187)
	No primaquine (N=253)	Very low dose total primaquine (<2 mg/kg)(N=1)	Low dose total primaquine (2 - <5 mg/kg)(N=92)	High dose total primaquine (≥ 5 mg/kg)(N=841)	
Age (years)					
Mean (SD)	22 (13)	23 (NA)	26 (15)	23 (14)	23 (14)
Age Category					
<5	17 (7%)	0 (0.0 %)	2 (2%)	40 (5%)	59 (5%)
5-<15	56 (22%)	0 (0.0 %)	19 (21%)	237 (28%)	312 (26%)
≥15	180 (71%)	1 (100%)	71 (77%)	564 (67%)	816 (69%)
Gender					
Male	167 (66%)	1 (100%)	71 (77%)	540 (64%)	779 (66%)
Female	86 (34%)	0 (0.0 %)	21 (23%)	301 (36%)	408 (34%)
Weight (kg)					
Mean (SD)	43 (15)	62 (NA)	52 (14)	43 (15)	43 (15)
Missing	1 (0.4%)	0 (0.0 %)	43 (46.7%)	0 (0.0 %)	44 (3.7%)
Malnutrition					
No	14 (6%)	0 (0.0 %)	2 (2%)	42 (5%)	58 (5%)
Yes	7 (3%)	0 (0.0 %)	1 (1%)	14 (2%)	22 (2%)
Missing	232 (91.7%)	1 (100%)	89 (96.7%)	785 (93.3%)	1107 (93.3%)
Fever day 0					
No	29 (11%)	0 (0.0 %)	24 (26%)	10 (1%)	63 (5%)
Yes	223 (88%)	1 (100%)	25 (27%)	831 (99%)	1080 (91%)
Missing	1 (0.4%)	0 (0.0 %)	43 (46.7%)	0 (0.0 %)	44 (3.7%)
P. vivax baseline parasitaemia					
Median (IQR)	5670 [1624, 11925]	12560 [12560, 12560]	4346 [1262, 7357]	4192 [1408, 12811]	4352 [1440, 12058]
Haemoglobin day 0 (g/dL)					
Mean (SD)	12 (1.7)	12 (NA)	13 (1.6)	12 (1.7)	12 (1.7)
Missing	4 (1.6%)	0 (0.0 %)	43 (46.7%)	6 (0.7%)	53 (4.5%)
PQ daily dose (mg/kg)					
Mean (SD)		0.48 (NA)	3.7 (0.50)	7.0 (0.60)	6.7 (1.2)
Duration of PQ treatment					
7 days		0 (0.0 %)	4 (4%)	323 (38%)	327 (35%)
14 days		1 (100%)	88 (96%)	518 (62%)	607 (65%)
Method to calculate PQ dose					
Per actual dose		1 (100%)	11 (12%)	840 (100%)	852 (91%)
Per dosing protocol		0 (0.0 %)	81 (88%)	1 (0%)	82 (9%)
Start day of PQ treatment					
Day 0		1 (100%)	15 (16%)	759 (90%)	775 (83%)
Day 1		0 (0.0 %)	76 (83%)	1 (0%)	77 (8%)
Day 2		0 (0.0 %)	1 (1%)	81 (10%)	82 (9%)

(continued)

	No primaquine (N=253)	Very low dose total primaquine (<2 mg/kg)(N=1)	Low dose total primaquine (2 - <5 mg/kg)(N=92)	High dose total primaquine (≥ 5 mg/kg)(N=841)	Total (N=1187)
Day 3		0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Day 4		0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Day 5		0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Day 6		0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Level of PQ supervision					
Unsupervised		0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Partially supervised		0 (0.0 %)	38 (41%)	1 (0%)	39 (4%)
Fully supervised		1 (100%)	54 (59%)	840 (100%)	895 (96%)
Was PQ taken with food?					
No		1 (100%)	3 (3%)	194 (23%)	198 (21%)
Yes		0 (0.0 %)	89 (97%)	647 (77%)	736 (79%)
Recommended		0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Other treatment given					
AL	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
AsAq	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
AsMf	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Cq	253 (100%)	1 (100%)	90 (98%)	519 (62%)	863 (73%)
DP	0 (0.0 %)	0 (0.0 %)	2 (2%)	322 (38%)	324 (27%)
Transmission intensity of the site (note 1)					
Low	31 (12%)	0 (0.0 %)	32 (35%)	1 (0%)	64 (5%)
Moderate	222 (88%)	1 (100%)	60 (65%)	840 (100%)	1123 (95%)
High	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Not available	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Geographical region					
Africa	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Americas	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Asia-Pacific	253 (100%)	1 (100%)	92 (100%)	841 (100%)	1187 (100%)
Relapse Peridocity					
Low periodicity	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
High periodicity	253 (100%)	1 (100%)	92 (100%)	841 (100%)	1187 (100%)
G6PD categories (Qualitative test)					
<30%	24 (9%)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	24 (2%)
≥30%	229 (91%)	0 (0.0 %)	92 (100%)	841 (100%)	1162 (98%)
Missing	0 (0.0 %)	1 (100%)	0 (0.0 %)	0 (0.0 %)	1 (0.1%)
G6PD categories (Quantitative test)					
<30%	24 (9%)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	24 (2%)
30-<70%	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
≥70%	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Missing	229 (90.5%)	1 (100%)	92 (100%)	841 (100%)	1163 (98.0%)

¹ Transmission intensity was classified as low if there was a *P. vivax* malaria incidence rate of <1 per 1000 persons, moderate if 1 to <10 per 1000 persons and high if ≥10 per 1000 persons.

1.3: Risk of recurrence

Kaplan-Meier survival analysis was used to calculate risk of recurrence between day 7 and 365. Patients were left censored at day 7 and right censored at the first of: the day last reviewed, the last day prior to a 60-day blood smear gap or the last day of study follow up. Outcomes were stratified by primaquine treatment arm: no primaquine, low total dose primaquine (2 to <5 mg/kg) and high total dose primaquine (≥ 5 mg/kg). Very low total dose primaquine (<2 mg/kg) was not presented due to low numbers of patients treated with this dose.




Figure 0_eff: Kaplan-Meier figure of cumulative risk of recurrence between day 7 and day 365 for primaquine treatment category. Please interpret the results of this figure with caution as there may not always be paired treatment comparisons in the original studies contributing to these pooled results.

Cox regression analysis for the time to first vivax recurrence between day 7 and 180 was performed to determine the effect of primaquine dose. Analysis was restricted to patients treated with daily primaquine or no primaquine. Potential confounders including sex, age and baseline parasitaemia were adjusted for with shared frailty for study site.

Similar but separate multivariable Cox regression analyses were undertaken to investigate primaquine duration, also adjusting for total actual mg/kg dose, in i) patients treated with low total dose primaquine and ii) patients treated with high total dose primaquine.

Care should be taken when interpreting these results, as model assumptions have not been fully assessed in this automated report format.




Figure 1_eff: Hazard ratio between day 7 and day 180 for A: total dose of primaquine and B: 14-day vs 7-day primaquine duration, stratified by total dose of primaquine

2: HAEMATOLOGY

2.1: Description

Haematological safety is a key concern for clinicians and policymakers in the implementation of primaquine radical cure, due to the risk of haemolysis in patients with G6PD deficiency. This individual patient data meta-analysis was conducted to assess the evidence for adverse haematological outcomes related to primaquine dose, with consideration of patients G6PD status.

Inclusion in the haematological safety meta-analysis was restricted to studies with 28 days or more follow up, patients with data on day 0 parasitaemia, patients with available data on day 0 haemoglobin levels or haematocrit, patients with an available haemoglobin measurement on at least one more day during the follow-up period and patients with data on daily primaquine dose.

The haematology study included 1219 patients across 8 study sites, from 6 studies.

2.2 Characteristics of Study Population

Table 1_saf: Characteristics of the study population for the safety study analysis, categorised by total primaquine category

	Primaquine Treatment				Total (N=1219)
	No primaquine (N=249)	Low dose daily primaquine (<0.375 mg/kg/day) (N=46)	Intermediate dose daily primaquine (≥ 0.375 & <0.75 mg/kg/day) (N=553)	High dose daily primaquine (≥ 0.75 mg/kg/day) (N=371)	
Age (years)					
Mean (SD)	22 (13)	25 (12)	23 (13)	23 (13)	23 (13)
Age Category					
<5	16 (6.43%)	1 (2.17%)	27 (4.88%)	13 (3.50%)	57 (4.68%)
5- <15	54 (21.69%)	5 (10.87%)	141 (25.50%)	94 (25.34%)	294 (24.12%)
≥ 15	179 (71.89%)	40 (86.96%)	385 (69.62%)	264 (71.16%)	868 (71.21%)
Gender					
Male	164 (65.86%)	41 (89.13%)	370 (66.91%)	253 (68.19%)	828 (67.92%)
Female	85 (34.14%)	5 (10.87%)	183 (33.09%)	118 (31.81%)	391 (32.08%)
Weight (kg)					
Mean (SD)	43 (15)	51 (15)	43 (15)	43 (15)	43 (15)
Missing	1 (0.4%)	0 (0.0 %)	43 (7.8%)	42 (11.3%)	86 (7.1%)
Malnutrition					
No	13 (5.22%)	1 (2.17%)	30 (5.42%)	12 (3.23%)	56 (4.59%)
Yes	7 (2.81%)	0 (0.00%)	10 (1.81%)	5 (1.35%)	22 (1.80%)
Missing	229 (92.0%)	45 (97.8%)	513 (92.8%)	354 (95.4%)	1141 (93.6%)
Fever day 0					
No	29 (11.65%)	24 (52.17%)	5 (0.90%)	5 (1.35%)	63 (5.17%)
Yes	219 (87.95%)	22 (47.83%)	548 (99.10%)	366 (98.65%)	1155 (94.75%)
Missing	1 (0.4%)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	1 (0.1%)
P. vivax baseline parasitaemia					
Median (IQR)	5536 [1616, 11304]	5675 [4068, 11994]	4256 [1344, 12811]	4898 [1664, 14318]	4679 [1536, 12874]
Haemoglobin day 0 (g/dL)					
Mean (SD)	12 (1.7)	12 (1.7)	12 (1.8)	12 (1.7)	12 (1.8)
PQ daily dose (mg/kg)					
Mean (SD)		4.1 (0.86)	6.7 (1.2)	7.0 (0.49)	6.7 (1.1)
Missing		0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0(0%)
Mean (SD)		14 (0)	13 (1.9)	7.1 (0.72)	11 (3.5)
Missing		0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	

(continued)

	No primaquine (N=249)	Low dose daily primaquine (<0.375 mg/kg/day) (N=46)	Intermediate dose daily primaquine (≥ 0.375 & <0.75 mg/kg/day) (N=553)	High dose daily primaquine (≥ 0.75 mg/kg/day) (N=371)	Total (N=1219)
Method to calculate PQ dose					
Per dosing protocol		38 (82.61%)	44 (7.96%)	42 (11.32%)	124 (12.78%)
Missing		0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0(0%)
Start day of PQ treatment					
Day 1		36 (78.26%)	1 (0.18%)	0 (0.00%)	37 (3.81%)
Day 2		0 (0.00%)	37 (6.69%)	45 (12.13%)	82 (8.45%)
Day 3		0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
Day 4		0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
Day 5		0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
Day 6		0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
Missing		0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0(0%)
Level of PQ supervision					
Partially supervised		38 (82.61%)	1 (0.18%)	0 (0.00%)	39 (4.02%)
Fully supervised		8 (17.39%)	552 (99.82%)	371 (100.00%)	931 (95.98%)
Was PQ taken with food?					
No		4 (8.70%)	187 (33.82%)	1 (0.27%)	192 (19.79%)
Recommended		0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
Other treatment given					
AL	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
AsAq	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
Cq	249 (100.00%)	44 (95.65%)	351 (63.47%)	166 (44.74%)	810 (66.45%)
DP	0 (0.00%)	2 (4.35%)	159 (28.75%)	163 (43.94%)	324 (26.58%)
Missing	0 (0.0 %)	0 (0.0 %)	43 (7.8%)	42 (11.3%)	85 (7.0%)
Transmission intensity of the site (note 1)					
Low	31 (12.45%)	32 (69.57%)	44 (7.96%)	42 (11.32%)	149 (12.22%)
Moderate	218 (87.55%)	14 (30.43%)	509 (92.04%)	329 (88.68%)	1070 (87.78%)
High	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
Not available	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
Geographical region					
Africa	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
Americas	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
Asia-Pacific	249 (100.00%)	46 (100.00%)	553 (100.00%)	371 (100.00%)	1219 (100.00%)
Relapse Periodicity					
Low periodicity	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
High periodicity	249 (100.00%)	46 (100.00%)	553 (100.00%)	371 (100.00%)	1219 (100.00%)
G6PD categories (Qualitative test)					
$<30\%$	24 (9.64%)	0 (0.00%)	0 (0.00%)	0 (0.00%)	24 (1.97%)
$\geq 30\%$	223 (89.56%)	46 (100.00%)	509 (92.04%)	329 (88.68%)	1107 (90.81%)
Unknown	2 (0.80%)	0 (0.00%)	44 (7.96%)	42 (11.32%)	88 (7.22%)
G6PD categories (Quantitative test)					
$<30\%$	24 (9.64%)	0 (0.00%)	0 (0.00%)	0 (0.00%)	24 (1.97%)
30- $<70\%$	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
$\geq 70\%$	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)
Unknown	225 (90.36%)	46 (100.00%)	553 (100.00%)	371 (100.00%)	1195 (98.03%)

¹ Transmission intensity was classified as low if there was a *P. vivax* malaria incidence rate of <1 per 1000 persons, moderate if 1 to <10 per 1000 persons and high if ≥ 10 per 1000 persons.

2.3 Summary of the haematology outcomes

Table 2 below provides a summary of the outcome experienced within each primaquine treatment arm for participants with G6PD activity $\geq 30\%$.

Table 2_saf: Summary of safety outcomes, categorised by total primaquine category


	Primaquine Treatment				
	No primaquine	Low dose daily primaquine (<0.375 mg/kg/day)	Intermediate dose daily primaquine (0.375 & <0.75 mg/kg/day)	High dose daily primaquine (0.75 mg/kg/day)	Total
Drop in haemoglobin of >25% AND Hb below 7 g/dL					
No	37 (16.4 %)	42 (91.3 %)	367 (66.5 %)	365 (98.4 %)	811 (67.9 %)
Yes	0 (0.0 %)	0 (0.0 %)	4 (0.7 %)	5 (1.3 %)	9 (0.8 %)
Missing	188 (83.6%)	4 (8.7%)	181 (32.8%)	1 (0.3%)	374 (31.3%)
Drop in haemoglobin of >5 g/dL from baseline between days 1-14					
No	37 (16.4 %)	42 (91.3 %)	365 (66.1 %)	367 (98.9 %)	811 (67.9 %)
Yes	0 (0.0 %)	0 (0.0 %)	6 (1.1 %)	3 (0.8 %)	9 (0.8 %)
Missing	188 (83.6%)	4 (8.7%)	181 (32.8%)	1 (0.3%)	374 (31.3%)
Drop in haemoglobin to <5 g/dL between days 1 and 14					
No	37 (16.4 %)	42 (91.3 %)	370 (67.0 %)	368 (99.2 %)	817 (68.4 %)
Yes	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	2 (0.5 %)	2 (0.2 %)
Missing	188 (83.6%)	4 (8.7%)	182 (33.0%)	1 (0.3%)	375 (31.4%)
Anaemia developed at days 2 or 3					
Nil (Hb: >=11 g/dL)	26 (11.6 %)	28 (60.9 %)	228 (41.3 %)	213 (57.4 %)	495 (41.5 %)
Mild (Hb: >=8 g/dL & <11 g/dL)	2 (0.9 %)	6 (13.0 %)	74 (13.4 %)	91 (24.5 %)	173 (14.5 %)
Moderate (Hb: >=5 g/dL & <8 g/dL)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	3 (0.8 %)	3 (0.3 %)
Severe (Hb <5 g/dL)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Missing	197 (87.6%)	12 (26.1%)	250 (45.3%)	64 (17.3%)	523 (43.8%)
Anaemia developed at days 5-7					
Nil (Hb: >=11 g/dL)	28 (12.4 %)	30 (65.2 %)	228 (41.3 %)	203 (54.7 %)	489 (41.0 %)
Mild (Hb: >=8 g/dL & <11 g/dL)	2 (0.9 %)	2 (4.3 %)	73 (13.2 %)	103 (27.8 %)	180 (15.1 %)
Moderate (Hb: >=5 g/dL & <8 g/dL)	0 (0.0 %)	0 (0.0 %)	3 (0.5 %)	2 (0.5 %)	5 (0.4 %)
Severe (Hb <5 g/dL)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Missing	195 (86.7%)	14 (30.4%)	248 (44.9%)	63 (17.0%)	520 (43.6%)
Change in haemoglobin on days 2-3 from day 0					
Mean (SD)	-0.345 (0.943)	-0.605 (1.14)	-0.862 (1.19)	-0.957 (1.08)	-0.872 (1.13)
Missing	194 (86.2%)	4 (8.7%)	187 (33.9%)	2 (0.5%)	387 (32.4%)
Change in haemoglobin on days 5-7 from day 0					
Mean (SD)	-0.0886 (0.721)	-0.307 (1.09)	-0.874 (1.35)	-1.04 (1.28)	-0.887 (1.31)
Missing	190 (84.4%)	7 (15.2%)	187 (33.9%)	2 (0.5%)	386 (32.3%)
Relative percentage (%) change in haemoglobin on days 2-3 from day 0					
Mean (SD)	2.60 (8.11)	4.38 (9.14)	6.54 (12.6)	7.61 (8.76)	6.77 (10.7)
Missing	194 (86.2%)	4 (8.7%)	187 (33.9%)	2 (0.5%)	387 (32.4%)
Relative percentage (%) change in haemoglobin on days 5-7 from day 0					
Mean (SD)	0.637 (5.83)	2.03 (9.33)	6.47 (14.0)	8.16 (10.3)	6.78 (12.1)
Missing	190 (84.4%)	7 (15.2%)	187 (33.9%)	2 (0.5%)	386 (32.3%)

2.4: Change in Haemoglobin (Hb) levels between primaquine treatment groups

The following figure provides the estimated change in haemoglobin from day 0 for different primaquine doses at at day 2/3 and days 5/7, adjusted for baseline haemoglobin, age, sex and day 0 parasitaemia and allowing for clustering by study site, in participants with $\geq 30\%$ G6PD activity.

Care should be taken when interpreting these results, as model assumptions have not been fully assessed in this automated report format.

A



B




Figure 1_saf: The covariate-adjusted estimated change in Hb between primaquine daily dose groups on (A) days 2-3 and (B) days 5-7, in patients with $\geq 30\%$ G6PD activity.

3: TOLERABILITY

3.1: Description

This individual patient data meta-analysis was conducted in order to understand the effect of primaquine dose on the gastrointestinal side effects.

Inclusion in the gastrointestinal tolerability meta-analysis was restricted to studies with 28 days or more followup, data from pre-specified symptom questionnaires (symptom checklist), patients with data on vivax parasite count at baseline, patients starting primaquine by day 2, patients not receiving intermittent primaquine (defined as primaquine administered weekly or monthly, rather than daily) and patients with data on daily primaquine dose.

The tolerability study included 1167 patients across 5 study sites, from 4 studies.

Characteristics of Study Population

Table 1_tol: Characteristics of the study population for the tolerability study analysis, categorised by total primaquine category

	Primaquine Treatment				Total (N=1167)
	No primaquine (N=222)	Low dose daily primaquine (<0.375 mg/kg/day) (N=16)	Intermediate dose daily primaquine (≥ 0.375 & <0.75 mg/kg/day) (N=558)	High dose daily primaquine (≥ 0.75 mg/kg/day) (N=371)	
Age (years)					
Mean (SD)	21 (13)	19 (13)	23 (13)	23 (13)	23 (13)
Age Category					
<5	17 (8%)	1 (6%)	28 (5%)	13 (4%)	59 (5%)
5- <15	56 (25%)	5 (31%)	142 (25%)	94 (25%)	297 (25%)
≥ 15	149 (67%)	10 (62%)	388 (70%)	264 (71%)	811 (69%)
Gender					
Male	144 (65%)	13 (81%)	373 (67%)	253 (68%)	783 (67%)
Female	78 (35%)	3 (19%)	185 (33%)	118 (32%)	384 (33%)
Weight (kg)					
Mean (SD)	41 (15)	41 (18)	43 (15)	43 (15)	42 (15)
Missing	1 (0.5%)	0 (0.0 %)	43 (7.7%)	42 (11.3%)	86 (7.4%)
Malnutrition					
No	14 (6%)	1 (6%)	31 (6%)	12 (3%)	58 (5%)
Yes	7 (3%)	0 (0.0 %)	10 (2%)	5 (1%)	22 (2%)
Missing	201 (90.5%)	15 (93.8%)	517 (92.7%)	354 (95.4%)	1087 (93.1%)
Fever day 0					
No	4 (2%)	4 (25%)	4 (1%)	5 (1%)	17 (1%)
Yes	217 (98%)	12 (75%)	554 (99%)	366 (99%)	1149 (98%)
Missing	1 (0.5%)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	1 (0.1%)
P. vivax baseline parasitaemia					
Median (IQR)	5096 [1618, 11075])	5124 [3950, 12482])	4224 [1348, 12811])	4898 [1664, 14318])	4608 [1504, 12811])
Haemoglobin day 0 (g/dL)					
Mean (SD)	12 (1.7)	12 (2.0)	12 (1.8)	12 (1.7)	12 (1.8)
Missing	4 (1.8%)	0 (0.0 %)	6 (1.1%)	0 (0.0 %)	10 (0.9%)
PQ daily dose (mg/kg)					
Mean (SD)		4.6 (1.0)	6.7 (1.2)	7.0 (0.49)	6.8 (1.0)
Duration of PQ treatment					
7 days		0 (0.0 %)	45 (8%)	367 (99%)	412 (44%)
14 days		16 (100%)	513 (92%)	4 (1%)	533 (56%)

(continued)

	No primaquine (N=222)	Low dose daily primaquine (<0.375 mg/kg/day) (N=16)	Intermediate dose daily primaquine (≥ 0.375 & <0.75 mg/kg/day) (N=558)	High dose daily primaquine (≥ 0.75 mg/kg/day) (N=371)	Total (N=1167)
Method to calculate PQ dose					
Per actual dose		8 (50%)	515 (92%)	329 (89%)	852 (90%)
Per dosing protocol		8 (50%)	43 (8%)	42 (11%)	93 (10%)
Start day of PQ treatment					
Day 0		8 (50%)	521 (93%)	326 (88%)	855 (90%)
Day 1		8 (50%)	0 (0.0 %)	0 (0.0 %)	8 (1%)
Day 2		0 (0.0 %)	37 (7%)	45 (12%)	82 (9%)
Day 3		0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Day 4		0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Day 5		0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Day 6		0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Level of PQ supervision					
Unsupervised		0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Partially supervised		8 (50%)	0 (0.0 %)	0 (0.0 %)	8 (1%)
Fully supervised		8 (50%)	558 (100%)	371 (100%)	937 (99%)
Was PQ taken with food?					
No		4 (25%)	193 (35%)	1 (0%)	198 (21%)
Yes		12 (75%)	365 (65%)	370 (100%)	747 (79%)
Recommended		0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Other treatment given					
AL	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
AsAq	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Cq	222 (100%)	14 (88%)	356 (64%)	166 (45%)	758 (65%)
DP	0 (0.0 %)	2 (12%)	159 (28%)	163 (44%)	324 (28%)
Missing	0 (0.0 %)	0 (0.0 %)	43 (7.7%)	42 (11.3%)	85 (7.3%)
Transmission intensity of the site (note 1)					
Low	0 (0.0 %)	2 (12%)	43 (8%)	42 (11%)	87 (7%)
Moderate	222 (100%)	14 (88%)	515 (92%)	329 (89%)	1080 (93%)
High	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Not available	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Geographical region					
Africa	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Americas	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Asia-Pacific	222 (100%)	16 (100%)	558 (100%)	371 (100%)	1167 (100%)
Relapse Periodicity					
Low periodicity	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
High periodicity	222 (100%)	16 (100%)	558 (100%)	371 (100%)	1167 (100%)
G6PD categories (Qualitative test)					
$<30\%$	24 (11%)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	24 (2%)
$\geq 30\%$	198 (89%)	16 (100%)	514 (92%)	329 (89%)	1057 (91%)
Missing	0 (0.0 %)	0 (0.0 %)	44 (7.9%)	42 (11.3%)	86 (7.4%)
G6PD categories (Quantitative test)					
$<30\%$	24 (11%)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	24 (2%)
$30-<70\%$	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
$\geq 70\%$	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Missing	198 (89.2%)	16 (100%)	558 (100%)	371 (100%)	1143 (97.9%)

¹ Transmission intensity was classified as low if there was a *P. vivax* malaria incidence rate of <1 per 1000 persons, moderate if 1 to <10 per 1000 persons and high if ≥ 10 per 1000 persons.

3.3 Summary of the gastrointestinal tolerability outcomes

The primary endpoint for this analysis was a composite indicator including the presence of vomiting or anorexia or diarrhoea on days 5-7 after enrolment.

Secondary endpoints for this analysis were:

- the presence of vomiting, nausea, anorexia, abdominal pain, diarrhoea or dizziness assessed separately on days 5-7¹
- the presence of the composite endpoint including vomiting or anorexia or diarrhoea on day 0, days 1-2 and days 5-7, assessed separately

¹Assessment of nausea, dizziness and abdominal pain was restricted to patients older than 5 years

Table 2 provides a summary of the outcome experienced within each Primaquine treatment arm.

Table 2_tol: Summary of gastrointestinal outcomes, categorised by total primaquine category

	Primaquine Treatment				Total
	No primaquine	Low dose daily primaquine (<0.375 mg/kg/day)	Intermediate dose daily primaquine (0.375 & <0.75 mg/kg/day)	High dose daily primaquine (0.75 mg/kg/day)	
Outcomes include participants of all ages					
	(N=222)	(N=16)	(N=558)	(N=371)	(N=1167)
Composite on day 0					
No	0 (0.0 %)	3 (37.5 %)	33 (76.7 %)	38 (90.5 %)	74 (79.6 %)
Yes	0 (0.0 %)	5 (62.5 %)	10 (23.3 %)	4 (9.5 %)	19 (20.4 %)
Missing	222 (100%)	8 (50.0%)	515 (92.3%)	329 (88.7%)	1074 (92.0%)
Composite between days 1-2					
No	0 (0.0 %)	0 (0.0 %)	40 (93.0 %)	39 (92.9 %)	79 (92.9 %)
Yes	0 (0.0 %)	0 (0.0 %)	3 (7.0 %)	3 (7.1 %)	6 (7.1 %)
Missing	222 (100%)	16 (100%)	515 (92.3%)	329 (88.7%)	1082 (92.7%)
Composite between days 5-7					
No	0 (0.0 %)	0 (0.0 %)	43 (100.0 %)	42 (100.0 %)	85 (100.0 %)
Yes	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Missing	222 (100%)	16 (100%)	515 (92.3%)	329 (88.7%)	1082 (92.7%)
Vomiting between days 5-7					
No	0 (0.0 %)	0 (0.0 %)	43 (100.0 %)	42 (100.0 %)	85 (100.0 %)
Yes	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Missing	222 (100%)	16 (100%)	515 (92.3%)	329 (88.7%)	1082 (92.7%)
Anorexia between days 5-7					
No	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Yes	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Missing	222 (100%)	16 (100%)	558 (100%)	371 (100%)	1167 (100%)
Diarrhoea between days 5-7					
No	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Yes	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Outcomes restricted to participants >5 years old					
Missing	222 (100%) (N=201)	16 (100%) (N=15)	558 (100%) (N=517)	371 (100%) (N=354)	1167 (100%) (N=1087)
Nausea between days 5-7*					
No	0 (0.0 %)	0 (0.0 %)	43 (100.0 %)	42 (100.0 %)	85 (100.0 %)
Yes	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Missing	201 (100%)	15 (100%)	474 (91.7%)	312 (88.1%)	1002 (92.2%)
Abdominal pain between days 5-7*					
No	0 (0.0 %)	0 (0.0 %)	43 (100.0 %)	41 (97.6 %)	84 (98.8 %)
Yes	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	1 (2.4 %)	1 (1.2 %)
Missing	201 (100%)	15 (100%)	474 (91.7%)	312 (88.1%)	1002 (92.2%)
Dizziness between days 5-7*					
No	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)

(continued)

	No primaquine	Low dose daily primaquine (<0.375 mg/kg/day)	Intermediate dose daily primaquine (0.375 & <0.75 mg/kg/day)	High dose daily primaquine (≥ 0.75 mg/kg/day)	Total
Yes	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)	0 (0.0 %)
Missing	201 (100%)	15 (100%)	517 (100%)	354 (100%)	1087 (100%)

Figure 1_tol: Distribution of primaquine daily dose by primaquine mg/kg daily dose category. Primaquine daily dose categories: Low: <0.375 mg/kg/day, Int (intermediate): ≥ 0.375 mg/kg/day and <0.750 mg/kg/day, and High: ≥ 0.750 mg/kg/day

3.4: Risk of gastrointestinal intolerance

The risk of gastrointestinal intolerance on days 5-7 was calculated from the number of patients reporting the composite outcome as a proportion of the total number of patients asked about each of the individual components of the composite; i.e. those asked about vomiting or anorexia or diarrhoea on any day between days 5-7. The 95% confidence intervals (CIs) for the risks were calculated as exact binomial CIs. The risks were stratified by primaquine daily dose categories




Figure 2_tol: Risk of gastrointestinal intolerance by symptoms. For each outcome the risk was estimated as the number of individuals experiencing the symptom as a proportion of the number of individuals asked about the symptom on any day between days 5-7. The confidence intervals (CIs) are exact binomial CIs.

3.4.1: Adjusted association between primaquine daily dose categories and gastrointestinal intolerance days 5-7

Covariate-adjusted estimated proportion of patients with gastrointestinal symptoms on days 5–7 cannot be presented as there were no records for any gastrointestinal intolerances (experienced or not experienced) for the reference group (no primaquine)

3.4.2: Risk of Acute Vomiting on days 0-2 and 3-13

The unadjusted risk of vomiting within an hour of primaquine administration (acute vomiting) was calculated on days 0-2 and days 3-13 for each primaquine dose group.

Table 3_tol: Risk of acute vomiting on days 0-2 and 3-13 by primaquine daily dose categories




Figure 3_tol: Risk of acute vomiting on days 0-2 and 3-13 by primaquine daily dose categories. The confidence intervals (CIs) are exact binomial CIs.