



Artemether-Lumefantrine Dose Impact Study Group Acknowledgements

WWARN thanks the following investigators who contributed individual patient data from published or unpublished clinical studies to the AL Dose Impact Study Group pooled analysis investigating:

- the mg/kg dosing range of the clinically-adopted ACT partner drug lumefantrine; and
 - the effects of lumefantrine mg/kg dosing on clinical outcome.
1. Abdulla S, Sagara I, Borrmann S, D'Allessandro U, Gonzalez R, Hamel M, et al. Efficacy and safety of artemether-lumefantrine dispersible tablets compared with crushed commercial tablets in African infants and children with uncomplicated malaria: a randomised, single-blind, multicentre trial. *Lancet* 2008. 372(9652): p. 1819-27.
 2. Achan J, Tibenderana JK, Kyabayinze D, Wabwire Mangen F, Kamya MR, Dorsey G, et al. Effectiveness of quinine versus artemether-lumefantrine for treating uncomplicated falciparum malaria in Ugandan children: randomised trial. *British Medical Journal* 2009. 339: p. b2763.
 3. Adam I, Salah MT, Eltahir HG, Elhassan AH, Elmardi KA, Malik EM. Dihydroartemisinin-piperaquine versus artemether-lumefantrine, in the treatment of uncomplicated Plasmodium falciparum malaria in central Sudan. *Annals of Tropical Medicine and Parasitology* 2010. 104(4): p. 319-26.
 4. Arinaitwe E, Sandison TG, Wanzira H, Kakuru A, Homsy J, Kalamya J, et al. Artemether-lumefantrine versus dihydroartemisinin-piperaquine for falciparum malaria: a longitudinal, randomized trial in young Ugandan children. *Clinical Infectious Diseases* 2009. 49(11): p. 1629-37.
 5. Barnes K, and Raman J, Efficacy of Artemether-Lumefantrine in Kwa Zulu Natal, South Africa, 2002.
 6. Bassat Q, González R, Machevo S, Nahum A, Lyimo J, Maiga H, et al. Similar efficacy and safety of artemether-lumefantrine (Coartem(R)) in African infants and children with uncomplicated falciparum malaria across different body weight ranges. *Malaria Journal* 2011. 10: p. 369.
 7. Bassat Q, Mulenga M, Tinto H, Piola P, Borrmann S, Menendez C, et al. Dihydroartemisinin-piperaquine and artemether-lumefantrine for treating uncomplicated malaria in African children: a randomised, non-inferiority trial. *PLoS One* 2009. 4(11): p. e7871.
 8. Bell DJ, Wootton D, Mukaka M, Montgomery J, Kayange N, Chimpeni P, et al. Measurement of adherence, drug concentrations and the effectiveness of artemether-lumefantrine, chlorproguanil-dapsone or sulphadoxine-pyrimethamine in the treatment of uncomplicated malaria in Malawi. *Malaria Journal* 2009. 8: p. 204.
 9. Borrmann S, Sasi P, Mwai L, Bashraheil M, Abdallah A, Muriithi S, et al. Declining responsiveness of Plasmodium falciparum infections to artemisinin-based combination treatments on the Kenyan coast. *PLoS One* 2011. 6(11): p. e26005.
 10. Bousema JT, Schneider P, Gouagna LC, Drakeley CJ, Tostmann A, Houben R, et al., Moderate effect of artemisinin-based combination therapy on transmission of Plasmodium falciparum. *Journal of Infectious Diseases*, 2006. 193(8): p. 1151-9.
 11. Bukirwa H, Yeka A, Kamya MR, Talisuna A, Banek K, Bakyaite N, et al, Artemisinin combination therapies for treatment of uncomplicated malaria in Uganda. *PLoS Clinical Trials* 2006. 1(1): p. e7.
 12. Carrara VI, Zwang J, Ashley EA, Price RN, Stepniewska K, Barends M, et al. Changes in the treatment responses to artesunate-mefloquine on the northwestern border of Thailand during 13 years of continuous deployment. *PLoS One* 2009. 4(2): p. e4551.
 13. Carrasquilla G, Baron C, Monsell EM, Cousin M, Walter V, Lefevre G, et al. Randomized, prospective, three-arm study to confirm the auditory safety and efficacy of artemether-lumefantrine in Colombian patients with uncomplicated Plasmodium falciparum malaria. *American Journal of Tropical Medicine Hygiene* 2012. 86(1): p. 75-83.
 14. Clark TD, Njama-Meya D, Nzarubara B, Maiteki-Sebuguzi C, Greenhouse B, Staedke SG, et al. Incidence of malaria and efficacy of combination antimalarial therapies over 4 years in an urban cohort of Ugandan

- children. *PLoS One* 2010. 5(7): p. e11759.
15. Denis MB, Tsuyuoka R, Lim P, Lindegardh N, Yi P, Top SN, et al. Efficacy of artemether-lumefantrine for the treatment of uncomplicated falciparum malaria in northwest Cambodia. *Tropical Medicine and International Health* 2006. 11(12): p. 1800-7.
 16. Dorsey G, Staedke S, Clark TD, Njama-Meya D, Nzarubara B, Maiteki-Sebuguzi C, et al. Combination therapy for uncomplicated falciparum malaria in Ugandan children: a randomized trial. *JAMA* 2007. 297(20): p. 2210-9.
 17. Epicentre, Paris, France, Efficacy of amodiaquine-artesunate and artemether-lumefantrine for the treatment of uncomplicated Plasmodium falciparum malaria in Nimba country, Liberia, 2009.
 18. Espie E, Lima A, Atua B, Dhorda M, Flevaud L, Sompwe EM, et al. Efficacy of fixed-dose combination artesunate-amodiaquine versus artemether-lumefantrine for uncomplicated childhood Plasmodium falciparum malaria in Democratic Republic of Congo: a randomized non-inferiority trial. *Malaria Journal* 2012. 11: p. 174.
 19. Falade C, Makanga M, Premji Z, Ortmann CE, Stockmeyer M and de Palacios PI. Efficacy and safety of artemether-lumefantrine (Coartem) tablets (six-dose regimen) in African infants and children with acute, uncomplicated falciparum malaria. *Transactions of the Royal Society of Tropical Medicine* 2005. 99(6): p. 459-67.
 20. Faucher JF, Aubouy A, Adeothy A, Cottrell G, Doritchamou J, Gourmel B, et al. Comparison of sulfadoxine-pyrimethamine, unsupervised artemether-lumefantrine, and unsupervised artesunate-amodiaquine fixed-dose formulation for uncomplicated plasmodium falciparum malaria in Benin: a randomized effectiveness noninferiority trial. *Journal of Infectious Diseases* 2009. 200(1): p. 57-65.
 21. Faye B, Ndiaye JL, Tine R, Sylla K, Gueye A, Lô AC, et al. A randomized trial of artesunate mefloquine versus artemether lumefantrine for the treatment of uncomplicated Plasmodium falciparum malaria in Senegalese children. *American Journal of Tropical Medicine and Hygiene* 2010. 82(1): p. 140-4.
 22. Faye B, Offianan AT, Ndiaye JL, Tine RC, Toure W, Djoman K, et al. Efficacy and tolerability of artesunate-amodiaquine (Camoquin plus) versus artemether-lumefantrine (Coartem) against uncomplicated Plasmodium falciparum malaria: multisite trial in Senegal and Ivory Coast. *Tropical Medicine and International Health* 2010. 15(5): p. 608-13.
 23. Gadalla NB, Adam I, Elzaki SE, Bashir S, Mukhtar I, Oguike M, et al. Increased pfmdr1 copy number and sequence polymorphisms in Plasmodium falciparum isolates from Sudanese malaria patients treated with artemether-lumefantrine. *Antimicrobial Agents and Chemotherapy* 2011. 55(11): p. 5408-11.
 24. Gaye O, Ndiaye JL and Tine R. Evaluation of the efficacy and tolerance of Artesunate-Mefloquine in adults with uncomplicated malaria in Senegal, 2010.
 25. Gaye O. Study on the efficacy of antimalarial drugs in the suburbs of Dakar in collaboration with the NMCP and the WHO, 2010.
 26. Grivoyannis A. In vivo selection of Plasmodium falciparum pfcr1 K76 and pfmdr1 alleles by Artemether-Lumefantrine in Mali, 2009.
 27. Guthmann JP, Cohuet S, Rigutto C, Fortes F, Saraiva N, Kiguli J, et al. High efficacy of two artemisinin-based combinations (artesunate + amodiaquine and artemether + lumefantrine) in Caala, Central Angola. *American Journal of Tropical Medicine and Hygiene* 2006. 75(1): p. 143-5.
 28. Hodel EM, Kabanywany AM, Malila A, Zanolari B, Mercier T, Beck HP, et al. Residual antimalarials in malaria patients from Tanzania--implications on drug efficacy assessment and spread of parasite resistance. *PLoS One* 2009. 4(12): p. e8184.
 29. Hwang J, Alemayehu BH, Hoos D, Melaku Z, Tekleyohannes SG, Teshe T, et al. In vivo efficacy of artemether-lumefantrine against uncomplicated Plasmodium falciparum malaria in Central Ethiopia. *Malaria Journal* 2011. 10: p. 209.
 30. Juma EA, Obonyo CO, Akhwale WS and Ogutu BR. A randomized, open-label, comparative efficacy trial of artemether-lumefantrine suspension versus artemether-lumefantrine tablets for treatment of uncomplicated Plasmodium falciparum malaria in children in western Kenya. *Malaria Journal* 2008. 7: p. 262.
 31. Juma EA. Efficacy of co-administered amodiaquine plus artesunate and artemether/lumefantrine for the treatment of uncomplicated falciparum malaria in children less than five years in different epidemiological settings in Kenya, 2004.
 32. Kamugisha E JS, Minde M, Kataraihya J, Kongola G, Kironde F and Swedberg G. Efficacy of artemether-lumefantrine in treatment of malaria among under-fives and prevalence of drug resistance

- markers in Igombe-Mwanza, north-western Tanzania. *Malaria Journal* 2012. 27.
33. Kamya MR, Yeka A, Bukirwa H, Lugemwa M, Rwakimari JB, Staedke SG, et al. Artemether-lumefantrine versus dihydroartemisinin-piperaquine for treatment of malaria: a randomized trial. *PLoS Clinical Trials* 2007. 2(5): p. e20.
 34. Karunajeewa HA, Mueller I, Senn M, Lin E, Law I, Gomorra PS, et al. A trial of combination antimalarial therapies in children from Papua New Guinea. *New England Journal of Medicine* 2008. 359(24): p. 2545-57.
 35. Laminou I. Randomized, double blind study comparing the therapeutic efficacy and safety of artemether-lumefantrine and artesunate amodiaquine in Gaya, Niger, 2011.
 36. Laminou I. Test d'efficacite de l'association Arthemeter lumefantrine dans le traitement du paludisme non complique dans les 3 sites sentinelles de surveillance de Gaya, Tessaoua et Agadez commune, 2008.
 37. Lefevre G, Looareesuwan S, Treeprasertsuk S, Krudsood S, Silachamroon U, Gathmann I, et al. A clinical and pharmacokinetic trial of six doses of artemether-lumefantrine for multidrug-resistant *Plasmodium falciparum* malaria in Thailand. *American Journal of Tropical Medicine Hygiene* 2001. 64(5-6): p. 247-56.
 38. Martensson A, Ngasala B, Ursing J, Isabel Veiga M, Wiklund L, Membi C, et al. Influence of consecutive-day blood sampling on polymerase chain reaction-adjusted parasitological cure rates in an antimalarial-drug trial conducted in Tanzania. *Journal of Infectious Diseases* 2007. 195(4): p. 597-601.
 39. Martensson A, Stromberg J, Sisowath C, Msellem MI, Gil JP, Montgomery SM, et al. Efficacy of artesunate plus amodiaquine versus that of artemether-lumefantrine for the treatment of uncomplicated childhood *Plasmodium falciparum* malaria in Zanzibar, Tanzania. *Clinical Infectious Diseases* 2005. 41(8): p. 1079-86.
 40. Mayxay M, Khanthavong M, Chanthongthip O, Imwong M, Pongvongsa T, Hongvanthong B, et al. Efficacy of artemether-lumefantrine, the nationally-recommended artemisinin combination for the treatment of uncomplicated falciparum malaria, in southern Laos. *Malaria Journal* 2012. 11(1): p. 184.
 41. Mayxay M, Khanthavong M, Lindegardh N, Keola S, Barends M, Pongvongsa T, et al. Randomized comparison of chloroquine plus sulfadoxine-pyrimethamine versus artesunate plus mefloquine versus artemether-lumefantrine in the treatment of uncomplicated falciparum malaria in the Lao People's Democratic Republic. *Clinical Infectious Diseases* 2004. 39(8): p. 1139-47.
 42. Menan H, Faye O, Same-Ekobo A, Oga AS, Faye B, Kiki Barro CP, et al. Comparative study of the efficacy and tolerability of dihydroartemisinin-piperaquine-trimethoprim versus artemether-lumefantrine in the treatment of uncomplicated *Plasmodium falciparum* malaria in Cameroon, Ivory Coast and Senegal. *Malaria Journal* 2011. 10: p. 185.
 43. Menan, H, Etude de l'efficacite et de la tolerance des associations Artesunate + Amodiaquine et Artemether + Lumefantrine dans le traitement du paludisme simple a *Plasmodium falciparum* dans trois sites sentinelles de la Cote D'Ivoire, 2012.
 44. Mens PF, Sawa P, van Amsterdam SM, Versteeg I, Omar SA, Schallig HD, et al. A randomized trial to monitor the efficacy and effectiveness by QT-NASBA of artemether-lumefantrine versus dihydroartemisinin-piperaquine for treatment and transmission control of uncomplicated *Plasmodium falciparum* malaria in western Kenya. *Malaria Journal* 2008. 7: p. 237.
 45. Mwesigwa J, Parikh S, McGee B, German P, Drysdale T, Kalyango JN, et al. Pharmacokinetics of artemether-lumefantrine and artesunate-amodiaquine in children in Kampala, Uganda. *Antimicrobial Agents and Chemotherapy* 2010. 54(1): p. 52-9.
 46. Ndiaye JL, Faye B, Gueye A, Tine R, Ndiaye D, Tchania C, et al. Repeated treatment of recurrent uncomplicated *Plasmodium falciparum* malaria in Senegal with fixed-dose artesunate plus amodiaquine versus fixed-dose artemether plus lumefantrine: a randomized, open-label trial. *Malaria Journal* 2011. 10: p. 237.
 47. Ndiaye JL, Randrianarivelojosa M, Sagara I, Brasseur P, Ndiaye I, Faye B, et al. Randomized, multicentre assessment of the efficacy and safety of ASAQ--a fixed-dose artesunate-amodiaquine combination therapy in the treatment of uncomplicated *Plasmodium falciparum* malaria. *Malaria Journal* 2009. 8: p. 125.
 48. Nikiema F and Zongo I. Evolution of therapeutic efficacies of artemisinin-based combination therapies (ASAQ and AL) for treatment of uncomplicated falciparum malaria in Burkina Faso during five years adoption as first-line treatments, 2010.
 49. Ngasala BE, Malmberg M, Carlsson AM, Ferreira PE, Petzold MG, Blessborn D, et al. Effectiveness of artemether-lumefantrine provided by community health workers in under-five children with uncomplicated malaria in rural Tanzania: an open label prospective study. *Malaria Journal* 2011. 10: p. 64.
 50. Offianan A, Assis SB, Coulibaly A, N'guessan LT, Ako AA, Kadjo FK, et al. Assessment of the efficacy of first-line

antimalarial drugs after 5 years of deployment by the National Malaria Control Programme in Côte d'Ivoire. *Open Access Journal of Clinical Trials* 2011. 3: p. 67-76.

51. Piola P, Fogg C, Bajunirwe F, Biraro S, Grandesso F, Ruzagira E, et al. Supervised versus unsupervised intake of six-dose artemether-lumefantrine for treatment of acute, uncomplicated *Plasmodium falciparum* malaria in Mbarara, Uganda: a randomised trial. *Lancet* 2005. 365(9469): p. 1467-73.
52. Premji Z, Umeh RE, Owusu-Agyei S, Esamai F, Ezedinachi EU, Oguche S, et al. Chlorproguanil-dapsone-artesunate versus artemether-lumefantrine: a randomized, double-blind phase III trial in African children and adolescents with uncomplicated *Plasmodium falciparum* malaria. *PLoS One* 2009. 4(8): p. e6682.
53. Price RN, Uhlemann AC, van Vugt M, Brockman A, Hutagalung R, Nair S, et al. Molecular and pharmacological determinants of the therapeutic response to artemether-lumefantrine in multidrug-resistant *Plasmodium falciparum* malaria. *Clinical Infectious Diseases* 2006. 42(11): p. 1570-7.
54. Ratcliff A, Siswantoro H, Kenangalem E, Maristela R, Wuwung RM, Laihadi F, et al. Two fixed-dose artemisinin combinations for drug-resistant falciparum and vivax malaria in Papua, Indonesia: an open-label randomised comparison. *Lancet* 2007. 369(9563): p. 757-65.
55. Sagara I, Diallo A, Kone M, Coulibaly M, Diawara SI, Guindo O, et al. A randomized trial of artesunate-mefloquine versus artemether-lumefantrine for treatment of uncomplicated *Plasmodium falciparum* malaria in Mali. *American Journal of Tropical Medicine Hygiene* 2008. 79(5): p. 655-61.
56. Sawa P, Shekalaghe SA, Drakeley CJ, Sutherland CJ, Mweresa CK, Baidjoe AY, et al. Malaria transmission after artemether-lumefantrine and dihydroartemisinin-piperazine: a randomized trial. *Journal of Infectious Diseases* 2013. 207(11): p. 1637-45.
57. Smithuis F, Kyaw MK, Phe O, Win T, Aung PP, Oo AP, et al. Effectiveness of five artemisinin combination regimens with or without primaquine in uncomplicated falciparum malaria: an open-label randomised trial. *Lancet Infectious Diseases* 2010. 10(10): p. 673-81.
58. Sutherland CJ, Ord R, Dunyo S, Jawara M, Drakeley CJ, Alexander N, et al. Reduction of malaria transmission to *Anopheles* mosquitoes with a six-dose regimen of co-artemether. *PLoS Medicine* 2005. 2(4): p. e92.
59. Syafruddin D, Asih PBS and Joshi AN. Evaluation of the parasite clearance day following treatment with artesunate-amodiaquine in subject with uncomplicated *Plasmodium falciparum* malaria in West Sumba District, East Nusa Tenggara Province, Indonesia, 2011.
60. The 4ABC Study Group. A head-to-head comparison of four artemisinin-based combinations for treating uncomplicated malaria in African children: a randomized trial. *PLoS Medicine* 2011;8(11):e1001119
61. Toure OA, Kouame MG, Didier YJ, Berenger AA, Djerea K, Genevieve GO, Penali LK. Artesunate/mefloquine paediatric formulation vs. artemether/lumefantrine for the treatment of uncomplicated *Plasmodium falciparum* in Anonkoua route, Cote d'Ivoire. *Tropical Medicine and International Health* 2011. 16(3): p. 290-7.
62. Toure OA, Penali LK, Yapi JD, Ako BA, Toure W, Djerea K, et al. A comparative, randomized clinical trial of artemisinin/naphthoquine twice daily one day versus artemether/lumefantrine six doses regimen in children and adults with uncomplicated falciparum malaria in Cote d'Ivoire. *Malaria Journal* 2009. 8: p. 148.
63. Ursing J, Kofoed PE, Rodrigues A, Blessborn D, Thoft-Nielsen R, Bjorkman A, et al. Similar efficacy and tolerability of double-dose chloroquine and artemether-lumefantrine for treatment of *Plasmodium falciparum* infection in Guinea-Bissau: a randomized trial. *Journal Infectious Diseases* 2011. 203(1): p. 109-16.
64. van den Broek I, Kitz C, Al Attas S, Libama F, Balasegaram M, Guthmann JP, et al. Efficacy of three artemisinin combination therapies for the treatment of uncomplicated *Plasmodium falciparum* malaria in the Republic of Congo. *Malaria Journal* 2006. 5: p. 113.
65. van den Broek I, Maung UA, Peters A, Liem L, Kamal M, Rahman M, et al. Efficacy of chloroquine + sulfadoxine-pyrimethamine, mefloquine + artesunate and artemether + lumefantrine combination therapies to treat *Plasmodium falciparum* malaria in the Chittagong Hill Tracts, Bangladesh. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 2005. 99(10): p. 727-35.
66. van Vugt M, Looreesuwan S, Wilairatana P, McGready R, Villegas L, Gathmann I et al. Artemether-lumefantrine for the treatment of multidrug-resistant falciparum malaria. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 2000. 94(5): p. 545-8.
67. van Vugt MV, Wilairatana P, Gemperli B, Gathmann I, Phaipun L, Brockman A, et al. Efficacy of six doses of artemether-lumefantrine (benflumetol) in multidrug-resistant *Plasmodium falciparum* malaria. *American Journal of Tropical Medicine and Hygiene* 1999. 60(6): p. 936-42.

68. Yavo W, Faye B, Kuete T, Djohan V, Oga SA, Kassi RR, et al. Multicentric assessment of the efficacy and tolerability of dihydroartemisinin-piperaquine compared to artemether-lumefantrine in the treatment of uncomplicated Plasmodium falciparum malaria in sub-Saharan Africa. *Malaria Journal* 2011. 10: p. 198.
69. Yeka A, Dorsey G, Kamya MR, Talisuna A, Lugeswa M, Rwakimari JB, et al. Artemether-lumefantrine versus dihydroartemisinin-piperaquine for treating uncomplicated malaria: a randomized trial to guide policy in Uganda. *PLoS One* 2008. 3(6): p. e2390.
70. Yeka A, Tibenderana J, Achan J, D'Alessandro U and Talisuna AO. Efficacy of quinine, artemether-lumefantrine and dihydroartemisinin-piperaquine as rescue treatment for uncomplicated malaria in Ugandan children. *PLoS One* 2013. 8(1): p. e53772.
71. Zongo I, Dorsey G, Rouamba N, Dokomajilar C, Sere Y, Rosenthal PJ, et al. Randomized comparison of amodiaquine plus sulfadoxine-pyrimethamine, artemether-lumefantrine, and dihydroartemisinin-piperaquine for the treatment of uncomplicated Plasmodium falciparum malaria in Burkina Faso. *Clinical Infectious Diseases* 2007. 45(11): p. 1453-61.
72. Zongo I, Dorsey G, Rouamba N, Tinto H, Dokomajilar C, Guiguemde RT, et al. Artemether-lumefantrine versus amodiaquine plus sulfadoxine-pyrimethamine for uncomplicated falciparum malaria in Burkina Faso: a randomised non-inferiority trial. *Lancet* 2007. 369(9560): p. 491-8.